



Lilly lowers the price of Zepbound® (tirzepatide) single-dose vials

December 1, 2025

Lilly continues efforts to expand access for people living with obesity by lowering the price of Zepbound single-dose vials

INDIANAPOLIS, Dec. 1, 2025 -- Eli Lilly and Company (NYSE: LLY) announced Zepbound® (tirzepatide) single-dose vials will now be available at lower prices on LillyDirect, the company's digital healthcare platform. This update builds on Lilly's ongoing efforts to lower out-of-pocket costs for people living with obesity in the United States. Last month, Lilly [announced lower prices for Zepbound](#) multi-dose pens, subject to approval by the Food and Drug Administration (FDA). By lowering prices of Zepbound single-dose vials, Lilly is offering patients greater access and optionality with their treatment options.

With a valid prescription, patients can access the starting dose of Zepbound single-dose vial (2.5 mg) for as low as \$299 per month. The 5 mg dose will be available at \$399 per month. All other approved doses are \$449 per month for patients in the Zepbound Self Pay Journey Program. ¹

"Far too many people who need obesity treatments still face cost and coverage barriers," said Ilya Yuffa, executive vice president and president, Lilly USA and Global Customer Capabilities. "Today's action underscores Lilly's commitment to improving access across the obesity care landscape. We will keep working to provide more options – expanding choices for delivery devices and creating new pathways for access – so more people can get the medicines they need."

Zepbound Self Pay Journey Program* via LillyDirect

Dose	New Price	Prior Price
2.5 mg	\$299	\$349
5 mg	\$399	\$499
7.5 mg	\$449	\$499
10 mg	\$449	\$499
12.5 mg	\$449	\$499
15 mg	\$449	\$499

*Regular prices (per month) when not purchased via the Zepbound Self-Pay Journey Program: \$599 for 7.5 mg, \$699 for 10 mg, \$849 for 12.5 mg, \$1,049 for 15 mg.

Zepbound is an injectable prescription medicine that may help adults with obesity, or some adults with overweight who also have weight-related medical problems, to lose excess body weight and keep the weight off. It may also help adults with moderate-to-severe obstructive sleep apnea (OSA) and obesity to improve their OSA. Zepbound contains tirzepatide and should not be used with other tirzepatide-containing products or any GLP-1 receptor agonist medicines. It is not known whether Zepbound is safe and effective for use in children.

Zepbound is also available in 2.5 mg, 5 mg, 7.5 mg, 10 mg, 12.5 mg, or 15 mg per 0.5 mL doses in a single-dose pen (autoinjector). The 2.5 mg is a starting dose and not an approved maintenance dose. The recommended maintenance doses are 5 mg, 10 mg, or 15 mg injected subcutaneously once per week for weight reduction and long-term maintenance. The recommended maintenance doses are 10 mg or 15 mg for OSA.

Additional resources and details are available on [Zepbound.lilly.com](https://www.zepbound.lilly.com). Visit [LillyDirect.lilly.com](https://www.lillydirect.lilly.com) to learn more about pricing, eligibility and delivery options.

About Lilly

Lilly is a medicine company turning science into healing to make life better for people around the world. We've been pioneering life-changing discoveries for nearly 150 years, and today our medicines help tens of millions of people across the globe. Harnessing the power of biotechnology, chemistry and genetic medicine, our scientists are urgently advancing new discoveries to solve some of the world's most significant health challenges: redefining diabetes care; treating obesity and curtailing its most devastating long-term effects; advancing the fight against Alzheimer's disease; providing solutions to some of the most debilitating immune system disorders; and transforming the most difficult-to-treat cancers into manageable diseases. With each step toward a healthier world, we're motivated by one thing: making life better for millions more people. That includes delivering innovative clinical trials that reflect the diversity of our world and working to ensure our medicines are accessible and affordable. To learn more, visit [Lilly.com](https://www.lilly.com) and [Lilly.com/news](https://www.lilly.com/news), or follow us on [Facebook](#), [Instagram](#), and [LinkedIn](#). P-LLY

Trademarks and Trade Names

All trademarks or trade names referred to in this standby statement are the property of the company, or, to the extent trademarks or trade names belonging to other companies are references in this press release, the property of their respective owners. Solely for convenience, the trademarks and trade names in this press release are referred to without the ® and ™ symbols, but such references should not be construed as any indicator that the company or, to the extent applicable, their respective owners will not assert, to the fullest extent under applicable law, the company's or their rights thereto. We do not intend the use or display of other companies' trademarks and trade names to imply a relationship with, or endorsement or sponsorship of us by, any other companies.

Cautionary Statement Regarding Forward-Looking Statements

This press release contains forward-looking statements (as that term is defined in the Private Securities Litigation Reform Act of 1995), including statements about the supply treatment presentation, and access of Zepbound (tirzepatide) as a treatment for adults with obesity or overweight and reflects Lilly's current belief and expectations. However, as with any pharmaceutical product, there are substantial risks and uncertainties in the

process of drug research, development, and commercialization. Among other things, there can be no guarantee that future study results will be consistent with the results to date, that Zepbound will receive additional regulatory approvals, or that Lilly will execute its strategy as planned. For further discussion of these and other risks and uncertainties, see Lilly's most recent Form 10-K and Form 10-Q filings with the United States Securities and Exchange Commission. Except as required by law, Lilly undertakes no duty to update forward-looking statements to reflect events after the date of this release.

¹ Zepbound Self Pay Journey Program lowers the price of the 7.5 mg, 10 mg, 12.5 mg and 15 mg vials only on first fill and when prescriptions are refilled within 45 days of their previous delivery. The 12.5 mg and 15 mg vials are priced at \$849 and \$1,049, respectively, if refilled outside of the 45-day window.

Refer to: Courtney Kasinger; ckasinger@lilly.com; 317-501-7056 (Media)

Michael Czapar; czapar_michael_c@lilly.com; 317-617-0983 (Investors)

INDICATIONS AND SAFETY SUMMARY WITH WARNINGS

Zepbound® (ZEHP-bownd) is an injectable prescription medicine that may help adults with:

- obesity, or some adults with overweight who also have weight-related medical problems to lose excess body weight and keep the weight off.
- moderate-to-severe obstructive sleep apnea (OSA) and obesity to improve their OSA.

It should be used with a reduced-calorie diet and increased physical activity.

Zepbound contains tirzepatide and should not be used with other tirzepatide-containing products or any GLP-1 receptor agonist medicines. It is not known if Zepbound is safe and effective for use in children.

Warnings - Zepbound may cause tumors in the thyroid, including thyroid cancer. Watch for possible symptoms, such as a lump or swelling in the neck, hoarseness, trouble swallowing, or shortness of breath. If you have any of these symptoms, tell your healthcare provider.

- Do not use Zepbound if you or any of your family have ever had a type of thyroid cancer called medullary thyroid carcinoma (MTC).
- Do not use Zepbound if you have Multiple Endocrine Neoplasia syndrome type 2 (MEN 2).
- Do not use Zepbound if you have had a serious allergic reaction to tirzepatide or any of the ingredients in Zepbound.

Zepbound may cause serious side effects, including:

Severe stomach problems. Stomach problems, sometimes severe, have been reported in people who use Zepbound. Tell your healthcare provider if you have stomach problems that are severe or will not go away.

Dehydration leading to kidney problems. Diarrhea, nausea, and vomiting may cause a loss of fluids (dehydration), which may cause kidney problems. It is important for you to drink fluids to help reduce your chance of dehydration.

Gallbladder problems. Gallbladder problems have happened in some people who use Zepbound. Tell your healthcare provider right away if you get symptoms of gallbladder problems, which may include pain in your upper stomach (abdomen), fever, yellowing of skin or eyes (jaundice), or clay-colored stools.

Inflammation of the pancreas (pancreatitis). Stop using Zepbound and call your healthcare provider right away if you have severe pain in your stomach area (abdomen) that will not go away, with or without vomiting. You may feel the pain from your abdomen to your back.

Serious allergic reactions. Stop using Zepbound and get medical help right away if you have any symptoms of a serious allergic reaction, including swelling of your face, lips, tongue or throat, problems breathing or swallowing, severe rash or itching, fainting or feeling dizzy, or very rapid heartbeat.

Low blood sugar (hypoglycemia). Your risk for getting low blood sugar may be higher if you use Zepbound with medicines that can cause low blood sugar, such as a sulfonylurea or insulin. **Signs and symptoms of low blood sugar** may include dizziness or light-headedness, sweating, confusion or drowsiness, headache, blurred vision, slurred speech, shakiness, fast heartbeat, anxiety, irritability, mood changes, hunger, weakness or feeling jittery.

Changes in vision in patients with type 2 diabetes. Tell your healthcare provider if you have changes in vision during treatment with Zepbound.

Depression or thoughts of suicide. You should pay attention to changes in your mood, behaviors, feelings or thoughts. Call your healthcare provider right away if you have any mental changes that are new, worse, or worry you.

Food or liquid getting into the lungs during surgery or other procedures that use anesthesia or deep sleepiness (deep sedation). Zepbound may increase the chance of food getting into your lungs during surgery or other procedures. Tell all your healthcare providers that you are taking Zepbound before you are scheduled to have surgery or other procedures.

Common side effects

The most common side effects of Zepbound include nausea, diarrhea, vomiting, constipation, stomach (abdominal) pain, indigestion, injection site reactions, feeling tired, allergic reactions, belching, hair loss, and heartburn. These are not all the possible side effects of Zepbound. Talk to your healthcare provider about any side effect that bothers you or doesn't go away.

Tell your doctor if you have any side effects. **You can report side effects at 1-800-FDA-1088 or www.fda.gov/medwatch.**

Before using Zepbound

- **Your healthcare provider should show you how to use Zepbound before you use it for the first time.**
- **Talk to your healthcare provider about low blood sugar and how to manage it. Tell your healthcare provider if you are taking medicines to treat diabetes including an insulin or sulfonylurea.**
- **If you take birth control pills by mouth, talk to your healthcare provider before you use Zepbound. Birth control pills may not work as well while using Zepbound.** Your healthcare provider may recommend another type of birth control for 4 weeks after you start Zepbound and for 4 weeks after each increase in your dose of Zepbound.

Review these questions with your healthcare provider:

- Do you have other medical conditions, including problems with your pancreas, or severe problems with your stomach, such as slowed emptying of your stomach (gastroparesis) or problems digesting food?
- Do you take diabetes medicines, such as insulin or sulfonylureas?
- Do you have a history of diabetic retinopathy?
- Are you scheduled to have surgery or other procedures that use anesthesia or deep sleepiness (deep sedation)?
- Do you take any other prescription medicines or over-the-counter drugs, vitamins, or herbal supplements?
- Are you pregnant, plan to become pregnant, breastfeeding, or plan to breastfeed? Zepbound may harm your unborn baby. Tell your healthcare provider if you become pregnant while using Zepbound. Zepbound may pass into your breast milk. You should talk with your healthcare provider about the best way to feed your baby while using Zepbound.
- **Pregnancy Exposure Registry:** There will be a pregnancy exposure registry for women who have taken Zepbound during pregnancy. The purpose of this registry is to collect information about the health of you and your baby. Talk to your healthcare provider about how you can take part in this registry, or you may contact Lilly at 1-800-LillyRx (1-800-545-5979).

How to take

- Read the Instructions for Use that come with Zepbound.
- Use Zepbound exactly as your healthcare provider says.
- Use Zepbound with a reduced-calorie diet and increased physical activity.
- Inject Zepbound under the skin (subcutaneously) of your stomach (abdomen), thigh, or have another person inject in the back of the upper arm. **Do not** inject ZEPBOUND into a muscle (intramuscularly) or vein (intravenously).
- **Use Zepbound 1 time each week, at any time of the day.**
- Change (rotate) your injection site with each weekly injection. **Do not** use the same site for each injection.
- If you take too much Zepbound, call your healthcare provider, call the Poison Help line at 1-800-222-1222 or go to the nearest hospital emergency room right away.

Zepbound injection is approved as a 2.5 mg, 5 mg, 7.5 mg, 10 mg, 12.5 mg, or 15 mg per 0.5 mL in single-dose pen or single-dose vial.

Learn more

Zepbound is a prescription medicine. For more information, call 1-800-LillyRx (1-800-545-5979) or go to www.zepbound.lilly.com.

This summary provides basic information about Zepbound but does not include all information known about this medicine. Read the information that comes with your prescription each time your prescription is filled. This information does not take the place of talking with your healthcare provider. Be sure to talk to your healthcare provider about Zepbound and how to take it. Your healthcare provider is the best person to help you decide if Zepbound is right for you.

ZP CON BS 25SEP2025

Zepbound® and its delivery device base are registered trademarks owned or licensed by Eli Lilly and Company, its subsidiaries, or affiliates.